Childhood Tuberculosis - Some Basic Issues

Jeffrey R. Starke, M.D. Baylor College of Medicine



TUBERCULOSIS IS A SOCIAL DISEASE WITH MEDICAL **IMPLICATIONS**







THE GREAT PARADOX OF TUBERCULOSIS – A CAUTIONARY TALE

By the use of drugs and BCG vaccines, we can

- cure tuberculosis disease [for less than 75 US dollars]
- prevent progression of tuberculosis infection into disease
- prevent a significant proportion of lifethreatening childhood tuberculosis [by BCG vaccination]



Yet, tuberculosis remains one of the three greatest infectious disease menaces for humans, and our inability to control it is our biggest public health failure.

EPIDEMIOLOGY OF CHILDHOOD TUBERCULOSIS IN SELECTED HIGH-BURDEN COUNTRIES

Childhood

| | Chilanooa | | Case Rate | | |
|--------------|-----------|---------------|-----------|--------------|--|
| Country | Cases | % Total Cases | Children | Total | |
| Afghanistan | 17.540 | 25.3 | 189 | 324 | |
| Brazil | 23,520 | 20.7 | 47 | 66 | |
| China | 86,978 | 5.3 | 27 | 129 | |
| Pakistan | 61,905 | 25.3 | 103 | 172 | |
| South Africa | 35,449 | 16.1 | 237 | 501 | |
| Zimbabwe | 12,267 | 16.1 | 221 | 603 | |
| Range | | 2.7-25.3 | 15-237 | 66-603 | |
| Total | 659,397 | 9.6 | | | |

Cooo Doto



Baylor College of Medicine

BURDEN OF CHILDHOOD TUBERCULOSIS IN DEVELOPING COUNTRIES

- 40% of cases may be in children
- May account for up to 10% of children's hospital admissions
- May account for 10% or more of hospital deaths in children
- Annual rate of infection up to 3% or higher in children in many countries



ECONOMIC AND SOCIAL BURDEN OF TUBERCULOSIS FOR CHILDREN

- Direct treatment costs
- Inpatient or institutional treatment
- Transportation costs for care
- Lost earnings of the family
- Redirecting resources from other needs
- Withdrawal from school
- Creation of orphans [parents with TB/ HIV]
- Stigmatization and discrimination






Should Primary Tuberculosis in Children Continue to be Neglected?


KHK Hsu *J Pediatrics* 1956; 48: 501

"Primary tuberculosis is the fountainhead of tuberculous diseases. When acquired during childhood, it may develop into serious tuberculous diseases within a short period of time, or remain latent during childhood, only to be become reactivated in adult life."



TRANSITIONS IN TUBERCULOSIS





Tuberculosis: A Low Tech Disease Even In 2011

~ 400 BC: Hippocrates [H&P]



■ 1882: Koch [TB culture]



1895: Rontgen [x-ray]



■ 1907: Mantoux [TST]





■ Late 1990's: IGRAs





Baylor College of Medicine

HOW IS TUBERCULOSIS DIAGNOSED?

Adults - Mycobacterial-based diagnosis

- positive sputum AFB smear 60% 75%
- positive sputum culture 90%
- positive tuberculin skin test 80% [HIV < 50%]

Children

- positive sputum or gastric AFB smear 10%
- positive sputum or gastric culture 10% 40%
- positive tuberculin skin test 50% 80%





DIAGNOSIS OF TUBERCULOSIS

Even in developed countries, the "gold standard" for the diagnosis of tuberculosis in children is the triad of:

a positive TST



an abnormal CXR and/or physical exam



Children's Hospital

a history of recent contact to an infectious adult case of TB

ADVERSE EFFECTS IN CHILDREN OF NOT PERFORMING CONTACT TRACING

- Missed cases
- Misdiagnosis tuberculosis not recognized
- Diagnose cases later increased morbidity and mortality
- Missed drug resistance [need source case culture and drug susceptibility results]
- Missed prevention treating exposed or infected children





A Difficult Case

A 19 mo old child in Houston presents acutely to an OSH with a several day history of fever, agitation and lethargy, and a possible generalized seizure.

An emergent EVD is placed, he is transferred to TCH. CXR is WNL, no family hx of TB. He is started on broad spectrum antibiotics









A Difficult Case


- CSF taken directly from the brain was normal
- CSF taken from the spine was very abnormal, indicating meningitis
- All cultures for tuberculosis were negative
- The tuberculin skin test yielded 0 mm of induration, and the QuantiFERON was indeterminate
- The T.Spot TB IGRA was "wicked positive"
- 3 weeks later, the mother's boyfriend's sister
 [child's babysitter] was found to have pulmonary
 TB
- The child had a typical course for TB meningitis and improved slowly on anti-TB medications





TB Studies on Adults & Children

| Moda | lity Adult Studies * | Pedi Studies * | Δ | Ratio adult to pedi | Year of 1 st publication |
|------------------|----------------------|----------------------|-------|------------------------|--|
| Cultur Solid | re: 400 | 38 | -362 | 11x | 1967 |
| Cultur Liquid | | 13 | -235 | 19x | 1966 |
| MODS | 37 | 7 | -30 | 5x | 2000 |
| PPD/1 | ST 6399 | 2031 | -4368 | 3x | 1907 |
| IGRA | 344 | 87 | -257 | 4x | 1999 |
| Xpert | 7 | 1 | -6 | 7 x | 2010 |
| PCR, a | ıll 1831 | 187 | -1644 | 10 x | 1990 |



*: PubMed Queries



New TB Tests Are Not Applied to Children

| <u>Test</u> | # of published | | |
|------------------------|---------------------|--|--|
| | studies in children | | |
| FNA | 140 | | |
| Fluorescent microscopy | 1 | | |
| LED-FM | 0 | | |
| MODS | 3 | | |
| Line-probe assays | 1 | | |
| LAMP | 0 | | |
| GeneXpert | 1 | | |



A Fundamental Question

Much of the attention for HIV and malaria has focused on children, but this has not happened for tuberculosis:



Why not?

SOME REASONS WHY CHILDHOOD TUBERCULOSIS HAS BEEN NEGLECTED

- Difficulty confirming the diagnosis
- Children are rarely contagious [public health "dead end"]
- Perception from TB policy makers that treating adults is enough
- Government programs fail to address children
- Lack of family centered contact tracing
- Perceived lack of scientific study and scrutiny
- Misplaced faith in the BCG vaccines
- Lack of industry support
- Inadequate advocacy by pediatricians





The Good News

- Improved advocacy on the international level has increased attention on children and TB
- Increased research capacity is being developed in high burden countries
- Better organization by pediatric TB experts
- Recent international meetings in Stockholm and by the NIH have fostered enthusiasm and collaboration
- Inclusion of children in major research efforts





Ten Truths About Childhood Tuberculosis

- Much childhood tuberculosis is preventable with simple, inexpensive measures.
- Non-preventable childhood pulmonary tuberculosis can be found earlier when it is easier to treat.
- Finding and treating adults with tuberculosis is not sufficient for controlling childhood TB.
- 4. BCG vaccines alone cannot control childhood tuberculosis.
 - Infection control has been largely neglected in high burden areas.





Ten Truths About Childhood Tuberculosis

- 6. Chest x-ray is essential to diagnose pulmonary tuberculosis in children.
- Adequate tuberculosis control for children requires a robust public health system with contact tracing and interagency cooperation.
- 8. Children with tuberculosis infection are a huge reservoir for future adult tuberculosis cases.
- 9. Childhood tuberculosis really is a window into the overall effectiveness of tuberculosis control.
 - Additional basic and translational research into childhood tuberculosis are badly needed.





CHILDHOOD TUBERCULOSIS: THE HIDDEN EPIDEMIC

Donald Int J Tuberc Lung Dis 2004; 8:627

"The time has come for the hidden epidemic of childhood tuberculosis to emerge from the shadow of adult tuberculosis and be seen as a neglected child health problem of considerable proportions in precisely those communities that do not have the resources to deal with it adequately."



