


TUBERCULOSIS IN WASHINGTON

Key Points


- Tuberculosis (TB) disproportionately affects vulnerable populations in Washington State (WA).
- On average, 4 cases of TB disease are diagnosed in WA each week.
- An estimated 200,000 people in WA are infected with TB.
- The COVID-19 pandemic has had an impact on TB, as they are both infectious diseases that primarily attack the lungs and can present with similar symptoms.

Reported TB Cases

In 2021, 199 cases of TB disease were reported in WA. This represents a 22.09% increase from the 163 TB cases reported in 2020.




Since 2012, incidence rates of TB in WA have progressed downward overall, similar to the trend seen in the United States (U.S.) as a whole. The WA TB incidence rate in 2021 was 2.6 cases per 100,000 population. While overall decline in TB incidence is encouraging, the current level of progress in the elimination of TB remains insufficient to eliminate TB in the United States in this century. We must continue to find and treat cases of active TB disease and also test and treat latent TB infection to prevent progression to disease and turn TB elimination into a reality.



*U.S. data for 2021 will be available on March 24th


County-Level

In 2021, three Washington counties reported 10 or more TB cases. Together, these three counties accounted for 75.4% of the 199 cases counted in WA. Twenty-one WA counties did not report any TB cases in 2021.



Age and Sex at Birth

Acting separately and together, factors such as age and sex can lead to an increased risk of TB. From 2017-2021, males aged 65 and older experienced the highest TB rate (5.7 cases per 100,000 population) of all age-sex groups.



// Risk Factors



The greatest TB disease burden in WA continues to be among persons born outside of the U.S., where risk of exposure to TB is greater. From 2017-2021, WA residents born somewhere other than the U.S., its territories, insular areas, or minor outlying islands accounted for 78.8% of all TB cases.



Known TB exposure is a risk factor for TB in WA. Approximately 200,000 WA residents are estimated to be infected with TB. From 2017-2021, 6.8% of TB cases were a known contact of a person with infectious TB disease, and 2.7% of TB cases had a history of incomplete treatment for latent tuberculosis infection.



Across the U.S., TB continues to affect those experiencing homelessness. In WA, 4.5% of TB cases from 2017-2021 reported having experienced homelessness within the year prior to their TB diagnosis.



Certain medical conditions and health behaviors lessen the ability of the immune system to keep TB infection from progressing to TB disease. Diabetes mellitus was the most frequently reported medical risk (23.1%) among all TB disease cases diagnosed from 2017-2021. Excess alcohol use was the most commonly recorded risk behavior (5.4%).



// Drug Resistance

Drug resistance is a concern for successful treatment of TB cases. There have been between one and eight reported cases of multidrug-resistant TB (MDR-TB) in WA per year over the last five years. To date, there have been no reported cases of extensively drug-resistant TB (XDR-TB) in WA .



// Mortality

TB is preventable, treatable, and curable, yet it stands as the most deadly of all infectious diseases worldwide. From 2017-2021, there were between four and ten TB-related deaths per year in WA.



**INVEST TO END TB.
SAVE LIVES.**

- Learn more about TB at www.doh.wa.gov/TB
- For questions about TB, contact Washington State Department of Health Tuberculosis Program: TBServices@doh.wa.gov, 206-418-5500

